
TECHNICAL CARD



Omega Spawn S is a high quality supplementary maturation feed for shrimp breeders. This maturation diet is a replacement for fresh foods that are normally used in shrimp maturation, such as squid or polychaetes. Omega Spawn S will help reducing biosecurity risk by replacing fresh food.

- Containing all necessary nutrients for broodstock
- Preventing denaturation of essential nutrients
- Highly attractive as it resembles fresh food
- Quick and easy preparation: DIY concept
- Convenient application
- Easily formed in different shapes, such as worm shape

| | BIOOUSLOCK IEEU | | |
|---------------------|-----------------|--------------------|-------|
| NUTRIENT | VALUE | NUTRIENT | VALUE |
| Moisture (%) | 8 | Vitamin A (IU/kg) | 20000 |
| Protein (%) | 56 | Vitamin C (mg/kg) | 1200 |
| Lipids (%) | 12 | Vitamin D3 (IU/kg) | 2500 |
| Ash (%) | 13 | Vitamin E (IU/kg) | 700 |
| Fiber (%) | 0.9 | n-3 HUFA (mg/g) | 20 |
| Calcium (g/kg) | 2,0 | DHA (mg/g) | 10 |
| Sodium (g/kg) | 1,4 | EPA (mg/g) | 5 |
| Phosphorus (g/kg) | 1,6 | | |
| | | | |
| INSTRUCTION FOR USE | | | |

Broadstack food



INGREDIENTS

Fish meal, squid meal, krill meal, pea protein, micro-algae, spirulina, fish gelatin, soy lecithin, yeast autolysate

INSTRUCTION FOR USE

- Weigh the feed according to the dosage (6 parts feed + 4 parts water).
 - Example: For 10 kg of final product: mix 6 kg of Omega Spawn S with 4L of freshwater.
- Essential to use 25-30°C water.
- Mix for 5 to 10 mins until you obtain a dough-like and homogenous product.
- After mixing, leave it at room temperature for 30-60 minutes.

Feeding method:

- 2-3% of the broodstock biomass
- 20-30% of the daily quantity of fresh food can be replaced by Omega Spawn S
- Adjust feed amounts according to local conditions.

STORAGE CONDITIONS

Store in a dry and cool place (best at 4°C, max. 20°C) SHELF LIFE 2 year **AVAILABLE SIZES**

Powder

PACKAGING Buckets of 6.0 kg

